

A/H/

Requested Patent: WO9301165A2

Title:

PHENYLGLYCINE DERIVATIVES PHARMACEUTICAL COMPOSITIONS
CONTAINING THEM AND THEIR USE IN THERAPY ;

Abstracted Patent: US5610183 ;

Publication Date: 1997-03-11 ;

Inventor(s): OWENS ANDREW P (GB); WILLIAMS BRIAN J (GB) ;

Applicant(s): MERCK SHARP DOHME (GB) ;

Application Number: US19930167870 19931217 ;

Priority Number(s):

GB19910014886 19910710; GB19920007278 19920402; WO1992GB01241
19920708 ;

IPC Classification: A61K31/195 ; C07C255/03 ; C07C271/10 ; C07C233/00 ;

Equivalents:

CA2110725, DE69208089D, EP0593615 (WO9301165), B1, A3, JP6509090T

ABSTRACT:


PCT No. PCT/GB92/01241 Sec. 371 Date Dec. 17, 1993 Sec. 102(e) Date Dec. 17, 1993 PCT
Filed Jul. 8, 1992 PCT Pub. No. WO93/01165 PCT Pub. Date Jan. 21, 1993 Compounds of
Formula (I) and pharmaceutically acceptable salts thereof: (I) wherein Q is optionally
substituted phenyl or naphthyl; X and Y are each H, C1-6alkyl, C2-6alkenyl or X and Y are
together =O; Z is O or S; R1 is H, optionally substituted C1-6alkyl, optionally substituted phenyl
(C1-4alkyl), C2-6alkenyl, COC1-6alkylhalo, CORa, COORa, CONHRa, COC1-6alkylNRaRb or
CONRaC1-6alkylCONRaRb; R2 is substituted C1-6alkyl, optionally substituted phenyl(C1-4alkyl),
C2-6alkenyl, C1-6alkylhalo, CORa, COORa, CONHRa, COC1-6alkylNRaRb or
CONRaC1-6alkylCONRaRb; or R1 and R2 together form a chain (CH2)p optionally substituted
by oxo; R3 is H [or C1-6alkyl]; R4 is H, C1-6alkyl or phenyl; and R5 is optionally substituted
(CH2)qphenyl; are tachykinin antagonists useful for treating pain.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07C 271/16, 229/14, 237/08 C07C 255/22, 233/18, 251/40 A61K 31/135		A2	(11) International Publication Number: WO 93/01165 (43) International Publication Date: 21 January 1993 (21.01.93)
(21) International Application Number: PCT/GB92/01241 (22) International Filing Date: 8 July 1992 (08.07.92) (30) Priority data: 9114886.6 10 July 1991 (10.07.91) GB 9207278.4 2 April 1992 (02.04.92) GB (71) Applicant (for all designated States except US): MERCK SHARP & DOHME LIMITED [GB/GB]; Hertford Road, Hoddesdon, Hertfordshire EN11 9BU (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): OWENS, Andrew, Pate [GB/GB]; 77 Dingle Road, Rushden, Northamptonshire NN10 9UE (GB). WILLIAMS, Brian, John [GB/GB]; 74 Godfrey Way, Great Dunmow, Essex CM6 2SE (GB).		(74) Agent: BARRETT-MAJOR, Julie, Diane; Merck & Co., Inc., European Patent Department, Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). (81) Designated States: CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, MC, NL, SE). Published Without international search report and to be republished upon receipt of that report.	

(54) Title: AROMATIC COMPOUNDS, COMPOSITIONS CONTAINING THEM AND THEIR USE IN THERAPY



(I)

(57) Abstract

Compounds of formula (I) and salts and prodrugs thereof wherein, Q is optionally substituted phenyl, heteroaryl or naphthyl; X and Y are each H, C₁₋₆ alkyl, C₂₋₆ alkenyl or X and Y are together =O; Z is O or S; R¹ is H, optionally substituted C₁₋₆ alkyl, optionally substituted phenyl(C₁₋₄ alkyl), C₂₋₆ alkylene, COC₁₋₆ alkylhalo, COR^a, COOR^a, CONHR^a, COC₁₋₆alkylNR^aR^b or CONR^aC₁₋₆alkylCONR^aR^b; R² is substituted C₁₋₆ alkyl, optionally substituted phenyl(C₁₋₄ alkyl), C₂₋₆ alkylene, C₁₋₆ alkylhalo, COR^a, COOR^a, CONHR^a, COC₁₋₆alkylNR^aR^b or CONR^aC₁₋₆alkylCONR^aR^b; or R¹ and R² together form in (CH₂)_p optionally substituted by oxo; R³ is H or C₁₋₆ alkyl; R⁴ is H, C₁₋₆ alkyl or phenyl; and R⁵ is optionally substituted (CH₂)_qphenyl; are tachykinin antagonists useful for treating pain.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MI	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

- 1 -

AROMATIC COMPOUNDS, COMPOSITIONS CONTAINING THEM AND
THEIR USE IN THERAPY

5 This invention relates to a class of aromatic compounds, which are useful as tachykinin receptor antagonists.

 The tachykinins are a group of naturally-occurring peptides found widely distributed throughout mammalian
10 tissues, both within the central nervous system and in the peripheral nervous and circulatory systems. The structures of three known mammalian tachykinins are as follows:

Substance P:

15 Arg-Pro-Lys-Pro-Gln-Gln-Phe-Phe-Gly-Leu-Met-NH₂

Neurokinin A:

His-Lys-Thr-Asp-Ser-Phe-Val-Gly-Leu-Met-NH₂

Neurokinin B:

Asp-Met-His-Asp-Phe-Phe-Val-Gly-Leu-Met-NH₂

20 Substance P is believed inter alia to be involved in the neurotransmission of pain sensations [Otsuka et al, "Role of Substance P as a Sensory Transmitter in Spinal Cord and Sympathetic Ganglia" in 1982 Substance P in the Nervous System, Ciba Foundation Symposium 91, 13-34
25 (published by Pitman) and Otsuka and Yanagisawa, "Does Substance P Act as a Pain Transmitter?" TIPS (Dec. 1987) 8 506-510], specifically in the transmission of pain in migraine (B.E.B. Sandberg et al, J. Med Chem, (1982) 25 1009) and in arthritis [Levine et al in Science (1984)
30 226 547-549]. These peptides have also been implicated in gastrointestinal (GI) disorders and diseases of the GI tract such as inflammatory bowel disease [Mantyh et al in Neuroscience (1988) 25 (3) 817-37 and D. Regoli in "Trends in Cluster Headache" Ed. Sicuteri et al Elsevier

- 2 -

Scientific Publishers, Amsterdam (1987) page 85)]. It is also hypothesised that there is a neurogenic mechanism for arthritis in which substance P may play a role [Kidd et al "A Neurogenic Mechanism for Symmetrical Arthritis" in The Lancet, 11 November 1989 and Grönblad et al "Neuropeptides in Synovium of Patients with Rheumatoid Arthritis and Osteoarthritis" in J. Rheumatol. (1988) 15(12) 1807-10]. Therefore, substance P is believed to be involved in the inflammatory response in diseases such as rheumatoid arthritis and osteoarthritis [O'Byrne et al in Arthritis and Rheumatism (1990) 33 1023-8]. Other disease areas where tachykinin antagonists are believed to be useful are allergic conditions [Hamelet et al Can. J. Pharmacol. Physiol. (1988) 66 1361-7], immunoregulation [Lotz et al Science (1988) 241 1218-21 and Kimball et al, J. Immunol. (1988) 141 (10) 3564-9], vasodilation, bronchospasm, reflex or neuronal control of the viscera [Mantyh et al, PNAS (1988) 85 3235-9] and, possibly by arresting or slowing β -amyloid-mediated neurodegenerative changes [Yankner et al, Science (1990) 250, 279-82], in senile dementia of the Alzheimer type, Alzheimer's disease and Down's Syndrome. Substance P may also play a role in demyelinating diseases such as multiple sclerosis and amyotrophic lateral sclerosis [J. Lubner-Narod et. al., poster to be presented at C.I.N.P. XVIIIth Congress, 28th June-2nd July, 1992, in press], and in disorders of bladder function such as bladder detrusor hyper-reflexia (Lancet, 16th May, 1992, 1239).

It has furthermore been suggested that tachykinins have utility in the following disorders: depression, dysthymic disorders, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina and Reynaud's disease, fibrosing and collagen diseases such as


- 3 -

scleroderma and eosinophilic fasciitis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, neuropathy, neuralgia, disorders
 5 related to immune enhancement or suppression such as systemic lupus erythematosus (European patent application no. 0 436 334), ophthalmic disease such as conjunctivitis, vernal conjunctivitis, and the like, and cutaneous diseases such as contact dermatitis, atopic dermatitis,
 10 urticaria, and other eczematoid dermatitis (European patent application no. 0 394 989).

In view of their metabolic instability, peptide derivatives are likely to be of limited utility as therapeutic agents. It is for this reason that non-
 15 peptide tachykinin receptor antagonists are sought.

In essence, this invention provides a class of potent non-peptide tachykinin receptor antagonists. By virtue of their non-peptide nature, the compounds of the present invention do not suffer from the shortcomings, in
 20 terms of metabolic instability, of known peptide-based tachykinin receptor antagonists.

European patent application no. 0 194 464 discloses compounds of formula (A):



(A)

wherein:


R^1 is loweralkyl, arylloweralkyl or optionally substituted phenyl;

- 4 -

R^2 is inter alia phenyl;
 R^3 is inter alia H or loweralkyl;
R is inter alia arylloweralkyl; and
n is inter alia 1.

5 The compounds are said to have anticonvulsant properties.

German patent application no. 28 51 435 discloses compounds of formula (B):



(B)

wherein:

R is H or CH₃; and


20 R^1 is inter alia a loweralkyl group substituted by an optionally substituted phenyl group.

The compounds are said to be useful in heart disease, obesity and diabetes, through their effect on carbohydrate metabolism.

25 Canadian patent application no 2,029,338 discloses compounds of formula (C):

30

- 5 -




(C)

10 wherein

Ar is optionally substituted phenyl, 1- or 2-naphthyl or 5- or 6-membered heteroaryl;

m is inter alia zero;

15 Y is H, Ar'NHC, Ar'NHC, R-C, RCH₂ or



n is inter alia 1;


W is inter alia H or C₁₋₂₀alkyl; and

25 Z is inter alia R-CH₂, where R is inter alia optionally substituted phenyl.

The compounds are said to be ACAT inhibitors useful in lowering blood cholesterol levels.

30 Dutch patent application no. 8003601 discloses compounds of formula (D):

- 6 -



wherein

R^1 is C_{1-10} alkyl;

R^2 and R^3 are H or C_{1-10} alkyl;

R^4 is inter alia optionally substituted phenyl;

15 R^5 is inter alia optionally substituted phenyl;

n is inter alia zero;

m is inter alia 1;

p is inter alia 1; and

q is inter alia zero.


20 The compounds are said to have anti-spasmodic,
anaesthetic and analgesic activity.

European patent applicaiton no. 0 384 088 discloses
(+) 2-N,N-dimethylamino-1-(3,4,5-
trimethoxyphenyl)methyloxy)-2-phenylbutane as an opiate
25 receptor ligand of use in the treatment of
gastrointestinal disorders.

The present invention provides a compound of formula
(I), or a salt or prodrug thereof:

30

- 7 -



(I)

wherein

Q represents optionally substituted phenyl, optionally substituted heteroaryl or optionally substituted naphthyl;

15 X and Y each represent H, C₁₋₆alkyl, C₂₋₆alkenyl or X and Y together form a group =O;

Z represents O or S;

R¹ represents H; C₁₋₆ alkyl optionally substituted by hydroxy, cyano, COR^a, COOR^a, CONR^aR^b,
 20 COC₁₋₄alkylNR^aR^b, CONR^aC₁₋₄alkylCONR^aR^b or NR^aR^b, (where R^a and R^b each independently represent H, C₁₋₆ alkyl, phenyl optionally substituted by one or more of C₁₋₆alkyl, C₁₋₆alkoxy, halo or trifluoromethyl or phenyl(C₁₋₄alkyl) optionally substituted in the phenyl
 25 ring by one or more of C₁₋₆ alkyl, C₁₋₆ alkoxy, halo or trifluoromethyl); phenyl(C₁₋₄ alkyl), (optionally substituted in the phenyl ring by one or more of C₁₋₆ alkyl, C₁₋₆ alkoxy, halo or trifluoromethyl); C₂₋₆ alkylene; COR^a; COOR^a; CONHR^a; COC₁₋₆alkylhalo;
 30 COC₁₋₆alkylNR^aR^b; or CONR^aC₁₋₆alkylCONR^aR^b, where R^a and R^b are as previously defined;

R² represents C₁₋₆ alkyl substituted by hydroxy, cyano, COR^a, COOR^a, CONR^aR^b, COC₁₋₄alkylNR^aR^b, CONR^aC₁₋₄alkylCONR^aR^b or NR^aR^b, (where R^a and R^b are as

- 8 -

above defined); phenyl(C₁₋₄ alkyl), (optionally substituted by one or more of C₁₋₆ alkyl, C₁₋₆ alkoxy, halo and trifluoromethyl in the phenyl ring); C₂₋₆ alkylene; COR^a; COOR^a; CONHR^a; COC₁₋₆alkylhalo;
 5 COC₁₋₆alkylNR^aR^b; or CONR^aC₁₋₆alkylCONR^aR^b, where R^a and R^b are as previously defined;

or R¹ and R² together form a chain (CH₂)_p optionally substituted by oxo; where p is 4 or 5 and where one methylene group may optionally be replaced by an oxygen
 10 atom or a group NR^x, where R^x is H or C₁₋₆ alkyl;

R³ represents H or C₁₋₆alkyl;

R⁴ represents H, C₁₋₆ alkyl or phenyl (optionally substituted by one or more of C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halo, cyano, nitro, trifluoromethyl,
 15 trimethylsilyl, SR^c, SOR^c, SO₂R^c, OR^c, NR^cR^d, NR^cCOR^d, NR^cCOOR^d, COOR^c or CONR^cR^d, where R^c and R^d each independently represent H, C₁₋₆ alkyl, phenyl or trifluoromethyl); and

R⁵ represents (CH₂)_qphenyl, wherein q is 0, 1, 2 or
 20 3, which may optionally be substituted in the phenyl ring by one or more of C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, SR^c, SOR^c, SO₂R^c, OR^c, NR^cR^d, NR^cCOR^d, NR^cCOOR^d, COOR^c or CONR^cR^d, where R^c and R^d are as above defined.

25 The alkyl, alkenyl and alkynyl groups referred to with respect to any of the above formulae may represent straight, branched or cyclic groups or a combination thereof. Thus, for example, suitable alkyl groups include methyl, ethyl, n- or iso-propyl, n-, sec-, iso-
 30 or tert-butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, and cycloalkyl-alkyl groups such as cyclopropylmethyl; suitable alkenyl groups include vinyl and allyl; and suitable alkynyl groups include propargyl.

- 9 -

The term "halo" as used herein includes fluoro, chloro, bromo and iodo, especially chloro and fluoro.

Where Q represents optionally substituted phenyl, heteroaryl or naphthyl, suitable substituents include

5 C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, SR^C, SOR^C, SO₂R^C, OR^C, NR^CR^d, NR^CCOR^d, NR^CCOOR^d, COOR^C or CONR^CR^d, (where R^C and R^d are as above defined), such as C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halo, cyano, nitro,

10 trifluoromethyl, SCH₃, SOCH₃, SO₂CH₃, OR^b, NR^bR^C, NR^bCOR^C, NR^bCOOR^C, COOR^b or CONR^bR^C, where R^b and R^C are as above defined. One or more substituents may be present and each may be located at any available ring position.

15 Suitable values of the group Q include phenyl, indolyl, naphthyl, thiophenyl, furanyl, pyridyl, indazolyl, imidazolyl, tetrazolyl, oxazolyl, benzothiophenyl, benzofuranyl and benzimidazolyl.

Preferably Q is optionally substituted phenyl, indolyl, furanyl, thiophenyl or naphthyl. More

20 preferably Q represents optionally substituted phenyl.

Preferably X and Y each represents H.

Preferably Z represents O.

In one subgroup of compounds according to the

25 invention, R¹ represents H; C₁₋₆ alkyl optionally substituted by hydroxy, cyano, COR^a, COOR^a, CONR^aR^b, COC₁₋₄alkylNR^aR^b, CONR^aC₁₋₄alkylCONR^aR^b or NR^aR^b, (where R^a and R^b each are as previously defined); phenyl(C₁₋₄ alkyl) (optionally substituted by one or more of C₁₋₆ alkyl, C₁₋₆ alkoxy, halo and trifluoromethyl in the

30 phenyl ring); C₂₋₆ alkylene; COR^a; COOR^a; CONHR^a; COC₁₋₄alkylNR^aR^b; or CONR^aC₁₋₄alkylCONR^aR^b (where R^a and R^b are as previously defined) and R² represents C₁₋₆ alkyl substituted by hydroxy, cyano, COR^a, COOR^a,

- 10 -

CONR^aR^b, COC₁₋₄alkylNR^aR^b, CONR^aC₁₋₄alkylCONR^aR^b or NR^aR^b, (where R^a and R^b each are as previously defined); phenyl(C₁₋₄ alkyl) (optionally substituted by one or more of C₁₋₆ alkyl, C₁₋₆ alkoxy, halo and trifluoromethyl in the phenyl ring); C₂₋₆ alkylene; COR^a; COOR^a; CONHR^a; 5 COC₁₋₄alkylNR^aR^b; or CONR^aC₁₋₄alkylCONR^aR^b (where R^a and R^b are as previously defined), or R¹ and R² together form a chain (CH₂)_p optionally substituted by oxo where p is 4 or 5 and where one methylene group may optionally be 10 replaced by an oxygen atom or a group NR^x, where R^x is H or C₁₋₆ alkyl.

In a further subgroup of compounds according to the invention R¹ represents H, C₁₋₆alkyl, phenyl(C₁₋₄alkyl), COR¹⁶, COOR¹⁶ or CONHR¹⁶, where R¹⁶ is C₁₋₆alkyl or 15 phenyl, and R² represents phenyl(C₁₋₄alkyl), COR¹⁶, COOR¹⁶ or CONHR¹⁶, where R¹⁶ is as previously defined.

Suitable values for the group R² include C₁₋₆alkyl substituted by, for example, cyano, hydroxy, NH₂, CO₂C₁₋₆alkyl, COR^a, CONR^aR^b, CONR^aCH₂CONR^aR^b, especially 20 CONHCH₂CONH₂, CON(CH₃)CH₂CONH(CH₃) or CON(CH₃)CH₂CON(CH₃)₂, COC₁₋₄alkylNR^aR^b, especially COCH₂NR^aR^b, such as COCH₂NH₂, C₁₋₆ alkenyl, especially allyl and formyl, and chains such as (CH₂)₄, (CH₂)₅ and (CH₂)₂O(CH₂)₂.

Suitable values for R¹ include those mentioned above for R², and especially H and C₁₋₆alkyl, such as methyl, ethyl, propyl and cyclopropylmethyl. 25

In a preferred group of compounds according to the invention, R¹ represents H or methyl and R² represents 30 C₁₋₂alkyl substituted by a group selected from hydroxy, cyano, CHO, CO₂(C₁₋₆alkyl), CONR^aR^b and NR^aR^b. More preferably R¹ represents H and R² represents CH₂CONH₂.

Suitable values for the group R³ include H and methyl, preferably H.


SUBSTITUTE SHEET

- 11 -

Preferably R^4 represents H.

Suitably R^5 represents $(CH_2)_q$ phenyl where q is 0, 1 or 2 and the phenyl group is substituted. Suitable phenyl substituents include methyl, methoxy, nitro, cyano, halo and trifluoromethyl. Preferably R^5 represents a substituted phenyl group. More preferably R^5 represents 3,5-dimethylphenyl or 3,5-bistrifluoromethylphenyl.

A preferred sub-group of compounds according to the invention is represented by formula (Ia)



(Ia)

wherein R^1 , R^2 , R^3 and Z are as defined for formula (I) above;

R^{20} and R^{21} each independently represent H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, SR^C , SOR^C , SO_2R^C , OR^C , $NR^C R^d$, $NR^C COR^d$, $NR^C COOR^d$, $COOR^C$ or $CONR^C R^d$, where R^C and R^d are as above defined;

each R^{23} represents C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, SR^C , SOR^C , SO_2R^C , OR^C , $NR^C R^d$, $NR^C COR^d$, $NR^C COOR^d$, $COOR^C$ or $CONR^C R^d$, where R^C and R^d are as above defined, preferably halo, such as chloro;

n is 0, 1, 2 or 3, preferably 0;

and salts and prodrugs thereof.

Particularly preferred are compounds of formula (Ia) wherein R^{20} and R^{21} are other than H and are located in

SUBSTITUTE SHEET

- 12 -

the 3- and 5-positions. Preferably R^{20} and R^{21} each represent methyl or, more preferably, trifluoromethyl.

For use in medicine, the salts of the compounds of formula (I) will be non-toxic pharmaceutically acceptable salts. Other salts may, however, be useful in the preparation of the compounds according to the invention or of their non-toxic pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds of this invention include acid addition salts which may, for example, be formed by mixing a solution of the compound according to the invention with a solution of a pharmaceutically acceptable non-toxic acid such as hydrochloric acid, oxalic acid, fumaric acid, p-toluenesulphonic acid, maleic acid, succinic acid, acetic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid. Salts of amine groups may also comprise quaternary ammonium salts in which the amino nitrogen atom carries a suitable organic group such as an alkyl, alkenyl, alkynyl or aralkyl moiety. Thus, for example, when both R^1 and R^2 are other than hydrogen, the nitrogen atom to which they are attached may be further substituted to give a quaternary ammonium salt. Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include metal salts such as alkali metal salts, e.g. sodium or potassium salts; and alkaline earth metal salts, e.g. calcium or magnesium salts.

The present invention includes within its scope prodrugs of the compounds of formula (I) above. In general, such prodrugs will be functional derivatives of the compounds of formula (I) which are readily convertible in vivo into the required compound of formula (I). Conventional procedures for the selection and preparation of suitable prodrug derivatives are

- 13 -

described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

The compounds according to the invention may exist both as enantiomers and as diastereomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention.

The invention also provides pharmaceutical compositions comprising one or more compounds of this invention in association with a pharmaceutically acceptable carrier. Preferably these compositions are in unit dosage forms such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, or suppositories, for oral, parenteral or rectal administration. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a non-toxic pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the

- 14 -

advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinylpyrrolidone or gelatin.

The present invention further provides a process for the preparation of a pharmaceutical composition comprising a compound of formula (I), which process comprises bringing a compound of formula (I) into association with a pharmaceutically acceptable carrier or excipient.

The compounds of formula (I) are of value in the treatment of a wide variety of clinical conditions which are characterised by the presence of an excess of tachykinin, in particular substance P, activity. These

- 15 -

may include disorders of the central nervous system such as anxiety, depression, psychosis and schizophrenia; neurodegenerative disorders such as dementia, including senile dementia of the Alzheimer type, Alzheimer's disease and Down's syndrome; demyelinating diseases such as MS and ALS and other neuropathological disorders such as peripheral neuropathy, including diabetic and chemotherapy-induced neuropathy, and postherpetic and other neuralgias; respiratory diseases such as chronic obstructive airways disease, bronchopneumonia, bronchospasm and asthma; inflammatory diseases such as inflammatory bowel disease, psoriasis, fibrositis, osteoarthritis and rheumatoid arthritis; allergies such as eczema and rhinitis; hypersensitivity disorders such as poison ivy; ophthalmic diseases such as conjunctivitis, vernal conjunctivitis, and the like; cutaneous diseases such as contact dermatitis, atopic dermatitis, urticaria, and other eczematoid dermatitis; addiction disorders such as alcoholism; stress related somatic disorders; reflex sympathetic dystrophy such as shoulder/hand syndrome; dysthymic disorders; adverse immunological reactions such as rejection of transplanted tissues and disorders related to immune enhancement or suppression such as systemic lupus erythematosus; gastrointestinal (GI) disorders and diseases of the GI tract such as disorders associated with the neuronal control of viscera such as ulcerative colitis, Crohn's disease and incontinence; disorders of bladder function such as bladder detrusor hyper-reflexia; fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis; disorders of blood flow caused by vasodilation and vasospastic diseases such as angina, migraine and Reynaud's disease; and pain or nociception, for example, that attributable to or associated with any

- 16 -

of the foregoing conditions, especially the transmission of pain in migraine. For example, the compounds of formula (I) may suitably be used in the treatment of disorders of the central nervous system such as anxiety, psychosis and schizophrenia; neurodegenerative disorders such as senile dementia of the Alzheimer type, Alzheimer's disease and Down's syndrome; respiratory diseases such as bronchospasm and asthma; inflammatory diseases such as inflammatory bowel disease, osteoarthritis and rheumatoid arthritis; adverse immunological reactions such as rejection of transplanted tissues; gastrointestinal (GI) disorders and diseases of the GI tract such as disorders associated with the neuronal control of viscera such as ulcerative colitis, Crohn's disease and incontinence; disorders of blood flow caused by vasodilation; and pain or nociception, for example, that attributable to or associated with any of the foregoing conditions or the transmission of pain in migraine.

The compounds of formula (I) are particularly useful in the treatment of pain or nociception and/or inflammation and disorders associated therewith such as, for example, neuropathy, such as diabetic and chemotherapy-induced neuropathy, postherpetic and other neuralgias, asthma, osteoarthritis, rheumatoid arthritis and especially migraine.

The present invention further provides a compound of formula (I), for use in therapy.

In the treatment of conditions involving actions of tachykinins released physiologically in response to noxious or other stimuli, a suitable dosage level is about 0.001 to 50 mg/kg per day, preferably about 0.005 to 10 mg/kg per day, and especially about 0.005 to 5


- 17 -

mg/kg per day. The compounds may be administered on a regimen of 1 to 4 times per day, preferably once daily.

According to a further or alternative aspect, the present invention provides a method of treatment of a human or animal subject suffering from or susceptible to a condition characterised by the presence of an excess of tachykinin which method comprises administering to a human or animal subject in need of such treatment an effective amount of a compound of formula (I), or a salt or prodrug thereof.

The present invention also provides the use of a compound of formula (I), or a salt or prodrug thereof, for the manufacture of a medicament for the treatment of conditions characterised by the presence of an excess of tachykinins.

According to one general process (a) the compounds according to the invention may be prepared by reaction of a compound of formula (II)



(11)


wherein Q, R¹, R², R³, X, Y and Z are defined as for
formula (I), with a compound of formula HalCHR⁴R⁵, where
30 R⁴ and R⁵ are as defined for formula (I) and Hal is halo,
such as bromo, chloro or iodo, in the presence of a base.

The reaction is conveniently carried out in a suitable organic solvent, such as ether, for example, tetrahydrofuran.

- 18 -

Suitable bases of use in the reaction include alkali or alkaline earth metal hydrides, for example, sodium hydride.

According to an alternative process, (b), compounds of formula (I) wherein R^1 , R^2 and R^3 are H may be prepared from intermediates of formula (IIIA)



(III)

wherein Q, X, Y, Z, R^4 and R^5 are as defined for formula (I) and W represents NOH (IIIA) by reduction.

Suitable reducing agents of use in the reduction include borane, catalytic hydrogenation in the presence of a suitable catalyst, such as a noble metal catalyst, for example palladium, which may be supported, for example, on carbon, nickel, dissolving metal reduction, for example using an alkali metal, such as sodium, in an alcohol, such as ethanol, or sodium amalgam.


A preferred reducing agent is borane. The borane reduction is preferably effected at elevated temperature, such as about 105-110°C.

Compounds of formula (I) may also be prepared from other compounds of formula (I). Thus, for example, compounds of formula (I) wherein R^1 represents H may be reacted with an optionally substituted alkylating or an acylating agent to produce compounds of formula (I) wherein R^1 represent an optionally substituted alkyl or an acyl group. Suitable procedures will be readily apparent to one skilled in the art.

- 19 -

Conversely, compounds of formula (I) wherein R^1 represents, for example, an acyl or a benzyl group, may be converted to compounds of formula (I) wherein R^1 represents H by, for example, hydrolysis or catalytic
5 hydrogenation. Suitable reagents and conditions are described in the accompanying examples, or will be readily apparent to one skilled in the art of organic chemistry.

Compounds of formula (II) wherein Z is O and X and Y together represent a group =O are commercially available
10 or may be prepared, for example, from intermediates of formula (IV)



(IV)

wherein Q and R^3 are as above defined and Ph represents phenyl, by hydrolysis.

25 The reaction is conveniently effected by heating a solution of the compound of formula (IV) in concentrated hydrochloric acid at reflux.

Compounds of formula (II) wherein Z is O and X and Y are =O may also be prepared by conventional procedures
30 for the preparation of amino acids which are well documented and are described, for example, in Chemistry and Biochemistry of the Amino Acids, ed. G. C. Barrett, Chapman and Hall, 1985.


- 20 -

Compounds of formula (II) wherein Z is S may be prepared from the corresponding compounds of formula (II) wherein Z is O by treating the latter compound with Lawesson's reagent or phosphorus pentasulphide in a suitable solvent, e.g. pyridine, at ambient or elevated temperature, suitably at the reflux temperature of the chosen solvent.

Compounds of formula (II) wherein X and Y represent H may be prepared from the corresponding compounds of formula (II) wherein X and Y together represent =O, by reduction.

Suitable reducing agents include metal hydrides, such as lithium aluminium hydride. The reaction is conveniently effected in a suitable organic solvent, such as ether, for example, tetrahydrofuran, suitably at elevated temperature, such as the reflux temperature of the solvent.

Intermediates of formula (IV) may be prepared from compounds of formula (V)



(V)

wherein R³ is as defined for formula (I), by reaction with a compound of formula Q-Hal wherein Hal is halo, such as bromo, chloro or iodo, in the presence of a base.

Suitable bases of use in the reaction include metal hydroxides, for example, sodium hydroxide. The reaction is conveniently effected in a mixture of water and a suitable organic solvent, such as a hydrocarbon, for

- 21 -

example, toluene, in the presence of a phase transfer catalyst, such as benzyltrimethyl ammonium chloride.


Compounds of formula (V) are commercially available or may be prepared by procedures readily apparent to one skilled in the art.

Compounds of formula Q-Hal are commercially available or may be prepared by conventional procedures known to those skilled in the art.

Compounds of formula (IIIA) may be prepared from the corresponding compounds of formula (III) wherein W represents O (IIIB) by reaction with hydroxylamine or a suitable derivative thereof.

Conveniently the reaction will be carried out in an aqueous organic solvent such as, for example, aqueous methanol. Preferably the reaction mixture will be buffered to approximately pH 4 by addition of a suitable salt, such as, for example sodium citrate.

Compounds of formula (IIIB) may be prepared by reaction of a compound of formula (VI)



(VI)

wherein X, Y, Z, R⁴ and R⁵ are as defined for formula (I) and R⁶⁰ represents an alkyl group, with a compound of formula Q-COHal, where Hal represents halo such as chloro or bromo and Q is as above defined, by the method described by J. W. Labadie *et. al.*, J. Org. Chem. Soc., 1983, 48, 4634.

- 22 -

Intermediates of formula (VI) may be prepared by analogous methods to that described by W. C. Still, J. Am. Chem. Soc., 1978, 100, 1481.

5 Intermediates of formula (III) are novel compounds and form a further aspect of the present invention.

Where the above-described process for the preparation of the compounds according to the invention gives rise to mixtures of stereoisomers these isomers may, if desired, be separated, suitably by conventional techniques such as preparative chromatography.

10 The novel compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The novel compounds may, for example, be resolved into their component enantiomers by standard techniques, such as the formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-d-tartaric acid and/or (+)-di-p-toluoyl-l-tartaric acid followed by fractional crystallization and regeneration of the free base. The novel compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary.

25 During any of the above synthetic sequences it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene and P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

- 23 -

The following Examples illustrate the preparation of compounds according to the invention.

5

- 24 -

DESCRIPTION 1: 1-(3,5-Dimethylphenyl)methoxy)-(2S)-2-ammonium-2-phenylethane oxylate salt

5 a) L-2-Phenylglycinol (5g) and di-t-butylidicarbonate (9.4g) was stirred in a dichloromethane solution (30ml) at room temperature for 3 hours. The precipitate which formed was filtered to give N-t-butoxycarbonyl-L-2-phenylglycinol, 4g.

10 b) N-t-Butoxycarbonyl-L-2-phenylglycinol (2g, part (a), was dissolved in a mixture of dimethylformamide (2ml) and tetrahydrofuran (10ml). Sodium hydride (80%) was added to this over a 10 minute period with stirring under nitrogen at room temperature. After a further 15 min, 3,5-dimethylbenzyl
15 bromide was added and the reaction left to stir for 4 hours. To the solution was added water (102ml), and the product extracted into ethyl acetate. The organic phase was washed twice with water and dried (MgSO_4). The solvent was removed in vacuo and the residue chromatographed in a column containing silica gel eluting with 100% petrol (bp 60-70°C) and 5% ethyl acetate/95% petrol (bp 60-80°C), to give 1-(3,5-dimethylphenyl)methoxy-(2S)-2-t-butoxycarbonylamino-2-phenylethane, 0.9g, mp 46-49°C. ^1H NMR (360MHz, CDCl_3) δ
20 7.38-7.22 (5H, m), 6.90 (1H, s), 6.83 (2H, s), 5.28 (1H, bs), 4.84 (1H, bs), 4.46 (1H, d, $J_{\text{gem}} = 11.9\text{Hz}$, $\text{OCH}_A\text{H}_B\text{Ph}$), 4.38 (1H, d, $J_{\text{gem}} = 11.9\text{Hz}$, $\text{OCH}_A\text{H}_B\text{Ph}$), 3.69 (1H, dd, $J = 9.74\text{Hz}$ and 4.39Hz), 3.60 (1H, m), 2.28 (6H, s), 1.40 (9H, s). m/z (Cl^+) 356
25 (M+H). Found: C, 74.37; H, 8.03; N, 3.97. $\text{C}_{22}\text{H}_{29}\text{NO}_3$ requires C, 74.33; H, 8.22; N, 3.94%.

- 25 -

c) 1-(3,5-Dimethylphenyl)methyloxy (2S)-2-t-butoxycarbonylamino-2-phenylethane (0.2g; part (b)) was dissolved in trifluoroacetic acid for 10 mins then evaporated to dryness. To a solution of the residue dissolved in methanol was added oxalic acid (70mg). The solvent was removed in vacuo and the residue recrystallized from diethylether/petrol (bp 60-80°C) to give the 1-(3,5-dimethylphenyl)methyloxy-(2S)-2-ammonium-2-phenylethane oxalate salt, 0.114g, 135-137°C. ¹H NMR (360MHz, DMSO) δ 7.51-7.38 (5H, m), 6.90 (3H, s), 4.52 (1H, dd, J = 7.37Hz and 5.0Hz), 4.48 (2H, d, J = 1.8Hz), 3.76-3.66 (2H, m), 2.24 (6H, s). Found: C, 65.07; H, 6.61; N, 4.00. C₁₇H₂₁NO.1.1C₂H₂O₄ requires C, 65.07; H, 6.60; N, 3.95%.

EXAMPLE 1: L-1-((3,5-Bis(trifluoromethyl)phenyl)methyloxy)-2-(t-butoxycarbonylamino)-2-phenylethane

2-N-t-Butoxycarbonylamino-L-2-phenylglycinol (Description 1a, 1g) was alkylated with 3,5-bis(trifluoromethyl)benzyl bromide in a manner analogous to that described in Description 1b to give the title compound, mp 52-53°C, m/e (CI⁺) = 464 (M+H), (CI⁻) = 462 (M-H). Found: C, 56.99; H, 4.67; N, 3.05: C₂₂H₂₂F₆NO₃ requires: C, 57.02; H, 5.00; N, 3.02%.

DESCRIPTION 2: L-2-Ammonium-1-((3,5-bis(trifluoromethyl)phenyl)methyloxy)-2-phenylethane oxalate salt

L-1-((3,5-Bis(trifluoromethyl)phenyl)methyloxy)-2-t-butoxycarbonylamino-L-2-phenylglycinol (Example 1) was

- 26 -

deprotected in an analogous manner to that described in Description 1c to give the title compound, mp 84-90°C. ¹H NMR (MeOH d₄, 360MHz) δ 7.97 (2H, s), 7.89 (1H, s), 7.47-7.41 (5H, m), 4.8 (2H, AB J_{gem} = 12.7Hz), 4.61 (1H, t, J = 5.93Hz), 3.9 (2H, d, J = 5.27Hz). m/e (CI⁺) = 364. Found: C, 48.46; H, 3.75; N, 2.94. C₁₇H₁₅F₆NO.1.4(C₂H₂O₄) requires C, 48.60; H, 3.67; N, 2.86%.

10 EXAMPLE 2: 1-((3,5-Dimethylphenyl)methyloxy)-(2S)-2-(((Carbomethoxy)methyl)amino)-2-phenylethane

1-((3,5-Dimethylphenyl)methyloxy)-(2S)-2-t-butoxycarbonylamino-2-phenylethane (1.9g, Description 1b) was treated with trifluoroacetic acid. After 10 minutes the solvent was removed in vacuo and the residue partitioned between dichloromethane and 2M-NaOH. The organic phase was washed with water, dried (MgSO₄) and evaporated in vacuo. To a solution of the residue in tetrahydrofuran (20ml) was added methyl bromoacetate (0.51ml) and triethylamine (0.56g) and the solution heated to reflux for 16 hours. Ethyl acetate and water were added and the organic phase washed with water, saturated brine and dried (MgSO₄). The residue obtained after removal of the solvent in vacuo was chromatographed on silica gel eluting with ethyl acetate/petroleum ether bp 60-80°C (1:1) to give the title compound 1.6g as an oil. Found: C, 73.13; H, 7.75; N, 4.32. C₂₀H₂₅NO₃ requires C, 73.37; H, 7.70; N, 4.28%.

25 EXAMPLE 3: (2S)-2-(((Carboxamido)methyl)ammonium)-1-((3,5-dimethylphenyl)methyloxy)-2-phenylethane oxalate salt

- 27 -

The product of Example 2 (0.8g) was dissolved in methanol (20ml) saturated with ammonia at 0°C. After the solution had been stored at +5°C for 4 days in a sealed flask the solution was evaporated to dryness and chromatographed on silica gel eluting with ethyl acetate. A solution of the product and oxalic acid in methanol was evaporated to dryness and the residue crystallised from hot diethyl ether to give the title compound, mp = 86-89°C. Found: C, 60.10; H, 6.35; N, 6.64. $C_{19}H_{24}N_2O_2 \cdot (C_2H_2O_4)_{1.3}$ requires C, 60.41; H, 6.24; N, 6.52%.

10 EXAMPLE 4: (2S) - 2 - ((3, 5 -
Bis(trifluoromethyl)phenyl)methoxy) - 2 -
((carbomethoxy)methyl)ammonium) - 2-phenylethane oxalate
salt

15 The title compound was prepared from (2S)-2-amino-2-phenyl-1-((bis(trifluoromethyl)phenyl)methoxy)-2-phenylethane (Description 2) using an analogous procedure to that described in Example 2, mp 95-97°C. Found: C, 49.33; H, 3.82; N, 2.61. $C_{20}H_{19}F_6NO_3 \cdot (C_2H_2O_4)_{1.2}$ requires C, 49.51; H, 3.97; N, 2.58%.

20 EXAMPLE 5: (2S) - 1 - ((3, 5 -
Bis(trifluoromethyl)phenyl)methoxy) - 2 -
((carboxamido)methyl)ammonium) - 2-phenylethane oxalate salt

25 The title compound was prepared from the product of Example 4 using an analogous procedure to that described in Example 3, mp = 147-151°C. Found: C, 49.35; H, 4.07; N, 5.47. $C_{19}H_{18}F_6N_2O_2 \cdot C_2H_2O_4$ requires C, 49.42; H, 3.95; N, 5.49%.

- 28 -

EXAMPLE 6: 2 - A m m o n i u m - 1 - ((3 , 5 -
bis(trifluoromethyl)phenyl)methyloxy)-2-(4-chlorophenyl)ethane.
oxalate salt monohydrate

5 a) Potassium hydroxide (27.0g), lithium chloride (6.75g) and triethylbenzylammonium chloride (1.84g) were dissolved in dichloromethane (40ml) and 33% aqueous ammonia solution (45ml) and gaseous ammonia bubbled through this solution at 0°C for 5 minutes. A solution of p-chlorobenzaldehyde (11.2g) and chloroform (10.2ml) in dichloromethane (40ml) was added
10 over 1 hour whilst maintaining the temperature at 0°C and this mixture stirred under a continuous stream of ammonia at 0°C for 6 hours and at room temperature for a further 18 hours. Water (120ml) and dichloromethane (40ml) were added, the aqueous layer was further extracted with dichloromethane,
15 concentrated *in vacuo*, filtered and adjusted to pH 6-7 with concentrated hydrochloric acid. On cooling to 0°C a precipitate formed and was collected by filtration, washed with ethanol and water and dried to give α -(4-chlorophenyl)glycine, 4.2g.

20 b) The amino acid (part (a), 4.2g) was added portionwise to a refluxing suspension of lithium aluminium hydride (1.2g) in tetrahydrofuran (72ml) and the mixture heated under reflux for a further 6 hours. The solution was cooled to room temperature and 10% sodium hydroxide solution (1.92ml) added dropwise followed by water (2.4ml). A solution of di-*tert*-butyl dicarbonate
25 (7.0g) and 4-dimethylaminopyridine (96mg) in dichloromethane (38ml) was added and the resulting mixture heated under reflux for 6 hours then cooled to room temperature and filtered

- 29 -

through a pad of anhydrous sodium sulphate. The resulting crude mixture was purified by chromatography on silica gel to give 2-(N-t-butoxycarbonylamino)-2-(4-chlorophenyl)ethanol.

5 c) The product (part (b), 1.0g) was alkylated with 3,5-bis(trifluoromethyl)benzyl bromide followed by deprotection with trifluoroacetic acid in an analogous manner to that described in Example 2b and 2c respectively to give the title compound. Found: C, 45.19; H, 3.27; N, 2.61. $C_{17}H_{14}ClF_6NO$. $C_2H_2O_4 \cdot H_2O$ requires C, 45.12; H, 3.59; N, 2.78%.

10 EXAMPLE 7: 2-Ammonium-1-((3,5-bis(trifluoromethyl)phenyl)methyloxy)-2-(3-chlorophenyl)ethane oxalate salt

15 The title compound was prepared by a method analogous to that described in Example 6 using m-chlorobenzaldehyde as starting material. Found: C, 46.60; H, 3.37; N, 2.89. $C_{17}H_{14}ClF_6NO \cdot C_2H_2O_4$ requires: C, 46.28; H, 3.31; N, 2.87%.

20 EXAMPLE 8: 2-Ammonium-1-((3,5-bis(trifluoromethyl)phenyl)methyloxy)-2-(2-chlorophenyl)ethane oxalate salt

25 a) Sodium hydride (80% suspension in oil, 2.03g) was washed twice with petroleum ether and to this solid was added tetrahydrofuran (50ml) and dimethylformamide (3ml) followed by the slow addition of a solution of 3,5-bis(trifluoromethyl)benzyl alcohol (15g) in tetrahydrofuran (50ml). After the effervescence had subsided (30 minutes) a

- 30 -

solution of tri-n-butyltinmethylene iodide (25.2g) was added. The solution was heated to reflux for 2 hours, cooled to room temperature and quenched by careful addition of petroleum ether bp 60-80°C (500ml) and water (200ml). The organic phase was washed with water and dried (MgSO₄). After removal of the solvent in vacuo the residue was distilled under reduced pressure bp_{0.8} = 140°C to give tri-n-butyl-((3,5-bis(trifluoromethyl)phenyl) methyloxymethyl)tin. ¹H NMR (360MHz, CDCl₃) δ 7.78 (1H, s, aryl 4CH), 7.76 (2H, s, aryl 2,6-CH), 4.52 (2H, s, aryl-CH₂).

b) 2-Chlorobenzoyl chloride (1.27ml), tri-n-butyl-((3,5-bis(trifluoromethyl)phenyl)methyloxymethyl)tin (part (a), 6.2g) and benzylchlorobis(triphenylphosphine)palladium (II) (80mg) were dissolved in chloroform (10ml) and the solution heated to 70°C for 36 hours. On cooling diethyl ether and saturated aqueous potassium fluoride were added and this solution filtered through a pad of Hyflo. The organic layer was washed with water, dried (MgSO₄) and reduced in vacuo giving a residue which was purified by chromatography on silica gel to give 2'-chloro-2-((3,5-bis(trifluoromethyl)phenyl) methyloxy)acetophenone, ¹H NMR (360MHz, CDCl₃) δ 7.79 (3H, bs, bis(CF₃) aryl CH₂,4,6), 7.54-7.26 (4H, m, aryl), 4.75 (2H, s), 4.76 (2H, s).

c) The product of part (b) (1.05g), sodium acetate (1.06g) and hydroxylamine hydrochloride were dissolved in methanol (20ml) and the solution stirred for 24 hours. Water was added and the mixture extracted with diethyl ether. The organic layer was washed with water, dried (MgSO₄) and reduced in vacuo to give

- 31 -

2'-chloro-2-((3,5-bis(trifluoromethyl)phenyl)methyloxy)acetophenone oxime.

d) A solution of borane (10ml, 1M in tetrahydrofuran) was added to a solution of the product of part (c) in tetrahydrofuran and the mixture heated under reflux for 48 hours. 5N hydrochloric acid was added and refluxing continued for 2 hours. On cooling diethyl ether was added and the solution washed with 2N aqueous sodium hydroxide. The organic layer was dried (MgSO_4) and reduced in vacuo to give a residue which was dissolved in methanol and treated with hydrogen gas at 50 psi over a 10% palladium on carbon catalyst for 2 hours. This mixture was filtered, reduced in vacuo and the residue partitioned between 5N hydrochloric acid and diethyl ether. The ether phase was washed with aqueous potassium hydroxide and dried (MgSO_4). Oxalic acid (150mg) was added and the solution allowed to stand for 24 hours when the title compound was collected as a precipitate (200mg). Found: C, 46.81; H, 3.63; N, 2.64%; $\text{C}_{17}\text{H}_{14}\text{ClF}_6\text{NO}(\text{COOH})_2$ requires: C, 46.28; H, 3.31; N, 2.87%.

20

EXAMPLE 9 : 1 - A m m o n i u m - 2 - ((bis(trifluoromethyl)phenyl)methyloxy)-1-phenylpropane oxalate salt hemihydrate

a) To a cooled (-30°C) solution of N-t-butoxycarbonyl (α -phenyl glycine (40g) and triethylamine (110ml) in dimethylformamide (500ml) was added iso-butylchloroformate (34.4ml) at such a rate that the temperature of the solution remained below -20°C . After stirring for 15 minutes to the

- 32 -

solution was added N,O-dimethylhydroxylamine hydrochloride (52g) and CH₂Cl₂ (800ml). The solution was stirred at room temperature for 16 hours. Ethyl acetate was added and the solution was washed with 10% aqueous citric acid solution, water (3 times), saturated NaHCO₃ solution and dried (MgSO₄).
5 The solvent was removed in vacuo and the resultant solid washed with hexane to give N'-t-butoxycarbonyl (α -phenyl)glycine N,O-dimethylhydroxamate, 37g.

b) To a cooled (0°C) solution of the product of part (a) (1.0g) in tetrahydrofuran (10ml) was slowly added methyl magnesium bromide (6.8ml). After stirring the solution at room temperature for 2 hours 10% aqueous citric acid and ethyl acetate were added and the organic phase washed further with water (2 times) and dried (MgSO₄). The solvent was removed in vacuo and to a solution of the residue in ethanol was added
15 sodium borohydride (0.13g) for 2 hours. The solvent was removed in vacuo and a solution of the residue in CH₂Cl₂ was washed with water (3 times) and dried (MgSO₄) to give upon evaporation in vacuo 1-N-t-butoxycarbonylamino-1-phenylpropan-2-ol as a mixture of diastereomers.
20

c) The product of part (b) was alkylated with 3,5-bis(trifluoromethyl)benzyl bromide in a manner analogous to that described in Description 1b to give 2-((bis(trifluoromethyl)phenyl)methoxy)-1-(N-t-butoxycarbonylamino)-1-phenylpropane as a mixture (approx
25 3:1) of diastereomers. This material (0.56g) was dissolved in trifluoroacetic acid (20ml) for 10 minutes followed by

- 33 -

evaporation in vacuo. A solution of the residue in CH_2Cl_2 was washed with 2N-NaOH, water (2 times), saturated brine and dried (MgSO_4). After removal of the solvent in vacuo the residue was crystallized by addition of oxalic acid (0.84g) followed by recrystallization from diethyl ether/petroleum ether to give the title compound, mp 60-62°C. Found: C, 50.17; H, 4.20; N, 2.84. $\text{C}_{18}\text{H}_{17}\text{F}_6\text{NO} \cdot \text{C}_2\text{H}_2\text{O}_4 \cdot 0.5(\text{H}_2\text{O})$ requires: C, 50.43; H, 4.23; N, 2.94%.

10 EXAMPLE 10: L-1-((3,5-Bis(trifluoromethyl)phenyl)methyloxy)-2-((cyanomethyl)amino)-2-phenylethane

A solution of L-2-Amino-1-(3,5-bis(trifluoromethyl)phenyl)methyloxy-2-phenylethane (1.5g, Description 2, free base), bromoacetonitrile (0.5g) and triethylamine (0.4g) in tetrahydrofuran (10ml) was heated under reflux for 6 hours. After cooling to room temperature and evaporation in vacuo, a solution of the residue in ethyl acetate was washed with water (3 times), saturated brine and dried (MgSO_4). The solvent was removed in vacuo and the residue purified by silica gel chromatography (eluting with 20-30% ethyl acetate in petroleum ether bp 60-80°C) to give the title compound as a crystalline solid. Found: C, 56.91; H, 4.21; N, 6.97. $\text{C}_{19}\text{H}_{15}\text{F}_6\text{N}_2\text{O}$ requires: C, 56.72; H, 4.01; N, 6.96%.

25 EXAMPLE 11: L-1-((Bis(trifluoromethyl)phenyl)methyloxy)-2-N-formamido-2-phenylethane

L-2-Amino-1-((bis(trifluoromethyl)phenyl)methyloxy)-2-phenylethane (Description 2, free base, 1.2g) was treated with

- 34 -

formic acetic anhydride (prepared by heating acetic anhydride
0.5g and formic acid 0.19ml at 80°C for 15 minutes) for 16 hours
at room temperature. The solution was evaporated to dryness to
give an oil which crystallized on standing. This solid was
recrystallized from ethyl acetate/petroleum ether to give a
5 mixture (approx 6:1) of L-1-
((bis(trifluoromethyl)phenyl)methyloxy)-2-N-formamido-2-
phenylethane and L-1-((bis(trifluoromethyl)phenyl)methyloxy)-
2-N-acetamido-2-phenylethane mp 96-99°C. Found: C, 55.50; H,
10 4.07; N, 3.49: $C_{18}H_{15}NO_2F_6$ requires C, 55.25; H, 3.86; N,
3.58%. m/e (CI^+) 392 (M+H), (CI^-) = 390 (M-H).

- 35 -

The following examples illustrate pharmaceutical compositions according to the invention.

5 EXAMPLE 12A Tablets containing 1-25mg of compound

	<u>Amount mg</u>		
Compound of formula (I)	1.0	2.0	25.0
Microcrystalline cellulose	20.0	20.0	20.0
Modified food corn starch	20.0	20.0	20.0
10 Lactose	58.5	57.5	34.5
Magnesium Stearate	0.5	0.5	0.5

EXAMPLE 12B Tablets containing 26-100mg of compound

	<u>Amount mg</u>		
15 Compound of formula (I)	26.0	50.0	100.0
Microcrystalline cellulose	80.0	80.0	80.0
Modified food corn starch	80.0	80.0	80.0
Lactose	213.5	189.5	139.5
Magnesium Stearate	0.5	0.5	0.5
20 The compound of formula (I), cellulose, lactose and a portion of the corn starch are mixed and granulated with 10% corn starch paste. The resulting granulation is sieved, dried and blended with the remainder of the corn starch and the magnesium stearate. The resulting			
25 granulation is then compressed into tablets containing 1.0mg, 2.0mg, 25.0mg, 26.0mg, 50.0mg and 100mg of the active compound per tablet.			

EXAMPLE 13 Parenteral injection

	<u>Amount mg</u>
30 Compound of formula (I)	1 to 100mg
Citric Acid Monohydrate	0.75mg
Sodium Phosphate	4.5mg
Sodium Chloride	9mg

- 36 -

Water for injection to 1ml
The sodium phosphate, citric acid monohydrate and sodium chloride are dissolved in a portion of the water. The compound of formula (I) is dissolved or suspended in the solution and made up to volume.

5 EXAMPLE 14 Topical formulation

	<u>Amount mg</u>
Compound of formula (I)	1-10g
10 Emulsifying Wax	30g
Liquid paraffin	20g
White Soft Paraffin	to 100g

The white soft paraffin is heated until molten. The liquid paraffin and emulsifying wax are incorporated and stirred until dissolved. The compound of formula (I) is added and stirring continued until dispersed. The mixture is then cooled until solid.

SUBSTANCE P ANTAGONISM ASSAY

20

A. Receptor Expression in Monkey Kidney Cell Line (COS)

To express the cloned human neurokinin-1- receptor (NK1R) transiently in COS, the cDNA for the human NK1R was cloned into the expression vector pCDM9 which was derived from pCDM8 (INVITROGEN) by inserting the ampicillin resistance gene (nucleotide 1973 to 2964 from BLUESCRIPT SK+ (trademark, STRATAGENE, La Jolla, CA, USA)) into the Sac II site. Transfection of 20 ug of the plasmid DNA into 10 million COS cells was achieved by electroporation in 800 µl of transfection buffer (135 mM NaCl, 1.2 mM CaCl₂, 1.2 mM MgCl₂, 2.4 mM K₂HPO₄, 0.6 mM KH₂PO₄, 10 mM glucose, 10 mM N-2-hydroxyethyl-piperazine-N'-2-ethane sulphonic acid (HEPES) pH 7.4) at 260 V and 950 µF using the IBI GENEZAPPER (trademark IBI, New

- 37 -

Haven, CT, USA). The cells were incubated in 10% fetal calf serum, 2 mM glutamine, 100U/ml penicillin-streptomycin, and 90% DMEM media (GIBCO, Grand Island, NY, USA) in 5% CO₂ at 37°C for three days before the binding assay.

B. Stable Expression in Chinese Hamster Ovarian Cell Line

To establish a stable cell line expressing cloned human NK1R, the cDNA was subcloned into the vector pRCCMV (INVITROGEN). Transfection of 20 ug of the plasmid DNA into CHO cells was achieved by electroporation in 800 µl of transfection buffer supplemented with 0.625 mg/ml Herring sperm DNA at 300 V and 950 µF using the IBI GENEZAPPER (IBI). The transfected cells were incubated in CHO media [10% fetal calf serum, 100 U/ml penicillin-streptomycin, 2 mM glutamine, 1/500 hypoxanthine-thymidine (ATCC), 90% IMDM media (JRH BIOSCIENCES, Lenexa, KS, USA), 0.7 mg/ml G418 (GIBCO)] in 5% CO₂ at 37°C until colonies were visible. Each colony was separated and propagated. The cell clone with the highest number of human NK1R was selected for subsequent applications such as drug screening.

C. Assay Protocol using COS or CHO

The binding assay of human NK1R expressed in either COS or CHO cells is based on the use of ¹²⁵I-substance P (¹²⁵I-SP, from DU PONT, Boston, MA) as a radioactively labeled ligand which competes with unlabeled substance P or any other ligand for binding to the human NK1R. Monolayer cell cultures of COS or CHO were dissociated by the non-enzymatic solution (SPECIALTY MEDIA, Lavellette, NJ) and resuspended in appropriate volume of the binding buffer (50 mM Tris pH 7.5, 5 mM MnCl₂, 150 mM NaCl, 0.04

- 38 -

mg/ml bacitracin, 0.004 mg/ml leupeptin, 0.2 mg/ml BSA, 0.01 mM phosphoramidon) such that 200 μ l of the cell suspension would give rise to about 10,000 cpm of specific ^{125}I -SP binding (approximately 50,000 to 200,000 cells). In the binding assay, 200 μ l of cells were added to a tube containing 20 μ l of 1.5 to 2.5 nM of ^{125}I -SP and 20 μ l of unlabeled substance P or any other test compound. The tubes were incubated at 4°C or at room temperature for 1 hour with gentle shaking. The bound radioactivity was separated from unbound radioactivity by GF/C filter (BRANDEL, Gaithersburg, MD) which was pre-wetted with 0.1% polyethylenimine. The filter was washed with 3 ml of wash buffer (50 mM Tris pH 7.5, 5 mM MnCl_2 , 150 mM NaCl) three times and its radioactivity was determined by gamma counter.

The activation of phospholipase C by NK1R may also be measured in CHO cells expressing the human NK1R by determining the accumulation of inositol monophosphate which is a degradation product of IP_3 . CHO cells are seeded in 12-well plate at 250,000 cells per well. After incubating in CHO media for 4 days, cells are loaded with 5 μ Ci of ^3H -myoinositol in 1 ml of media per well by overnight incubation. The extracellular radioactivity is removed by washing with phosphate buffered saline. LiCl is added to the well at final concentration of 10 mM with or without the test compound, and incubation is continued at 37°C for 15 min. Substance P is added to the well at final concentration of 0.3 nM to activate the human NK1R. After 30 min of incubation at 37°C, the medium is removed and 0.1 N HCl is added. Each well is sonicated at 4°C and extracted with CHCl_3 /methanol (1:1). The aqueous phase is applied to a 1 ml Dowex AG 1X8 ion exchange column. The column is washed with 0.1 N formic acid followed by 0.025 M ammonium formate-0.1 N formic acid.

- 39 -

The inositol monophosphate is eluted with 0.2 M ammonium formate-0.1 N formic acid and quantitated by beta counter.

The data in Table 1 were obtained for compounds of formula (I):


TABLE 1
SUBSTANCE P ANTAGONISM RESULTS

10	<u>Compound of Ex #</u>	<u>IC₅₀ @ NK1R (nM)</u>
	1	400
15	2	140
	3	100
	4	45
20	5	6
	10	15
25	11	60

- 40 -

CLAIMS:

1. A compound of formula (I), or a salt or prodrug thereof:



(I)

wherein

Q represents optionally substituted phenyl, optionally substituted heteroaryl or optionally substituted naphthyl;

- 20 X and Y each represent H, C₁₋₆alkyl, C₂₋₆alkenyl or X and Y together form a group =O;

Z represents O or S;

- 25 R¹ represents H; C₁₋₆ alkyl optionally substituted by hydroxy, cyano, COR^a, COOR^a, CONR^aR^b, COC₁₋₄alkylNR^aR^b, CONR^aC₁₋₄alkylCONR^aR^b or NR^aR^b, (where R^a and R^b each independently represent H, C₁₋₆ alkyl, phenyl optionally substituted by one or more of C₁₋₆alkyl, C₁₋₆alkoxy, halo or trifluoromethyl or phenyl(C₁₋₄alkyl) optionally substituted in the phenyl ring by one or more of C₁₋₆ alkyl, C₁₋₆ alkoxy, halo or trifluoromethyl); phenyl(C₁₋₄ alkyl) (optionally substituted in the phenyl ring by one or more of C₁₋₆ alkyl, C₁₋₆ alkoxy, halo or trifluoromethyl); C₂₋₆ alkylene; COR^a; COOR^a; CONHR^a; COC₁₋₆alkylhalo;
- 30

- 41 -

$\text{COC}_{1-6}\text{alkylNR}^a\text{R}^b$; or $\text{CONR}^a\text{C}_{1-6}\text{alkylCONR}^a\text{R}^b$, where R^a and R^b are as previously defined;

R^2 represents C_{1-6} alkyl substituted by hydroxy, cyano, COR^a , COOR^a , CONR^aR^b , $\text{COC}_{1-4}\text{alkylNR}^a\text{R}^b$, $\text{CONR}^a\text{C}_{1-4}\text{alkylCONR}^a\text{R}^b$ or NR^aR^b , (where R^a and R^b are as above defined); phenyl(C_{1-4} alkyl) (optionally substituted by one or more of C_{1-6} alkyl, C_{1-6} alkoxy, halo and trifluoromethyl in the phenyl ring); C_{2-6} alkylene; COR^a ; COOR^a ; CONHR^a ; $\text{COC}_{1-6}\text{alkylhalo}$;
 10 $\text{COC}_{1-6}\text{alkylNR}^a\text{R}^b$; or $\text{CONR}^a\text{C}_{1-6}\text{alkylCONR}^a\text{R}^b$, where R^a and R^b are as previously defined;

or R^1 and R^2 together form a chain $(\text{CH}_2)_p$ optionally substituted by oxo; where p is 4 or 5 and where one methylene group may optionally be replaced by an oxygen
 15 atom or a group NR^x , where R^x is H or C_{1-6} alkyl;

R^3 represents H or C_{1-6} alkyl;

R^4 represents H, C_{1-6} alkyl or phenyl (optionally substituted by one or more of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, SR^c , SOR^c , SO_2R^c , OR^c , NR^cR^d , NR^cCOR^d ,
 20 NR^cCOOR^d , COOR^c or CONR^cR^d , where R^c and R^d each independently represent H, C_{1-6} alkyl, phenyl or trifluoromethyl); and

R^5 represents $(\text{CH}_2)_q\text{phenyl}$, wherein q is 0, 1, 2 or
 25 3, which may optionally be substituted in the phenyl ring by one or more of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, SR^c , SOR^c , SO_2R^c , OR^c , NR^cR^d , NR^cCOR^d , NR^cCOOR^d , COOR^c or CONR^cR^d , where R^c and R^d are as above defined.

30

2. A compound as claimed in claim 1 wherein R^1 represents H; C_{1-6} alkyl optionally substituted by hydroxy, cyano, COR^a , COOR^a , CONR^aR^b , $\text{COC}_{1-4}\text{alkylNR}^a\text{R}^b$, $\text{CONR}^a\text{C}_{1-4}\text{alkylCONR}^a\text{R}^b$ or NR^aR^b , (where R^a and R^b each are

- 42 -

as previously defined); phenyl(C₁₋₄ alkyl) (optionally substituted in the phenyl ring by one or more of C₁₋₆ alkyl, C₁₋₆ alkoxy, halo and trifluoromethyl); C₂₋₆ alkylene; COR^a; COOR^a; CONHR^a; COC₁₋₄alkylNR^aR^b; or
5 CONR^aC₁₋₄alkylCONR^aR^b (where R^a and R^b are as previously defined) and R² represents C₁₋₆ alkyl substituted by hydroxy, cyano, COR^a, COOR^a, CONR^aR^b, COC₁₋₄alkylNR^aR^b, CONR^aC₁₋₄alkylCONR^aR^b or NR^aR^b, (where R^a and R^b each are as previously defined); phenyl(C₁₋₄ alkyl) (optionally
10 substituted in the phenyl ring by one or more of C₁₋₆ alkyl, C₁₋₆ alkoxy, halo and trifluoromethyl); C₂₋₆ alkylene; COR^a; COOR^a; CONHR^a; COC₁₋₄alkylNR^aR^b; or CONR^aC₁₋₄alkylCONR^aR^b (where R^a and R^b are as previously defined), or R¹ and R² together form a chain (CH₂)_p
15 optionally substituted by oxo where p is 4 or 5 and where one methylene group may optionally be replaced by an oxygen atom or a group NR^x, where R^x is H or C₁₋₆ alkyl.

3. A compound as claimed in claim 1 or claim 2
20 wherein R¹ represents H, C₁₋₆alkyl, phenyl(C₁₋₄alkyl), COR¹⁶, COOR¹⁶ or CONHR¹⁶, where R¹⁶ is C₁₋₆alkyl or phenyl, and R² represents phenyl(C₁₋₄alkyl), COR¹⁶, COOR¹⁶ or CONHR¹⁶, where R¹⁶ is as previously defined.

4. A compound as claimed in any preceding claim
25 wherein R¹ is H or methyl and R² is C₁₋₂alkyl substituted by a group selected from hydroxy, cyano, CHO, CO₂(C₁₋₆alkyl), CONR^aR^b and NR^aR^b.

5. A compound as claimed in any preceding claim
30 wherein Q is unsubstituted or substituted phenyl; R⁴ is H; and R⁵ is substituted phenyl.

- 43 -

6. A compound as claimed in any preceeding claim wherein X and Y both represent H and Z represents O.


7. A compound as claimed in claim 1 selected from:
 5 L-1-((3,5-bis(trifluoromethyl)phenyl)methyloxy)-2-(t-butoxycarbonylamino)-2-phenylethane;
 1-((3,5-dimethylphenyl)methyloxy)-2(S)-2-(((carbomethoxy)methyl)amino)-2-phenylethane;
 (2S)-2-(((carboxamido)methyl)ammonium)-1-((3,5-
 10 dimethylphenyl)methyloxy)-2-phenylethane;
 (2S)-2-((3,5-bis(trifluoromethyl)phenyl)methyloxy)-2-(((carbomethoxy)methyl)ammonium)-2-phenylethane;
 (2S)-2-((3,5-bis(trifluoromethyl)phenyl)methyloxy)-2-(((carboxamido)methyl)ammonium)-2-phenylethane;
 15 L-1-((3,5-bis(trifluoromethyl)phenyl)methyloxy)-2-((cyanomethyl)amino)-2-phenylethane;
 L-1-((3,5-bis(trifluoromethyl)phenyl)methyloxy)-2-N-formamido-2-phenylethane;
 and salts and prodrugs thereof.

20

8. A compound as claimed in any preceding claim for use in therapy.

9. A process for the preparation of a compound as
 25 claimed in any one of claims 1 to 7, which process comprises:

(a) reaction of a compound of formula (II):




(II)

- 44 -

where Q, R¹, R², R³, X, Y and Z are as defined as for formula (I), with a compound of formula HalCHR⁴R⁵, where R⁴ and R⁵ are as defined for formula (I) and Hal is halo, in the presence of a base; or

5

(b) reduction of a compound of formula (III):




(III)

wherein Q, X, Y, Z, R⁴ and R⁵ are as defined for formula (I) and W represents NOH;
 and, if necessary or desired, converting the compound of formula (I) so prepared into another compound of formula (I), or a salt or prodrug thereof.

15

10. An intermediate of formula (III):

20



(III)

wherein Q, X, Y, Z, R⁴ and R⁵ are as defined for formula (I) and W represents NOH.

30

11. A pharmaceutical composition comprising a compound as claimed in any one of claims 1 to 7 in association with a pharmaceutically acceptable carrier.

- 45 -

12. A method for the treatment or prevention of a physiological disorder associated with an excess of tachykinins, which method comprises administration to a patient in need thereof of a tachykinin-reducing amount of a compound according to claim 1.

13. A method according to claim 11 for the treatment or prevention of pain or inflammation.

14. A method according to claim 11 for the treatment or prevention of migraine.

15. A method according to claim 11 for the treatment or prevention of postherpetic neuralgia.